




Courtesy of Tourism Queensland

Impacts and adaptation strategies for a variable and changing climate in the **MARANOA AND DISTRICTS REGION**



This summary describes the likely impacts of a variable and changing climate on the major primary industries of the Maranoa and Districts (MD) region including grazing, dairy, cropping and horticulture, and the potential adaptation strategies which can be implemented to minimise climate risks.

Regional Profile



Macintyre River, Goondiwindi, Queensland

Courtesy of Tourism Queensland

The Maranoa and Districts (MD) region covers an area of 102,730 km² stretching from the Carnarvon Ranges in the north, to the Queensland-New South Wales border in the south, and from Mungallala in the west to about 90 km east of Roma. The main towns include Roma, Mitchell and St George.

This region has a highly variable summer rainfall with significant rainfall recorded in high intensity thunderstorm events. Average historical annual rainfall for the region is approximately 553 mm ranging from 573 mm at Mitchell to 533 mm at St George. Temperatures averaged over Roma, Mitchell and St George range from an annual average minimum of 13°C to an annual average maximum of 28°C. The region includes a significant part of the Queensland Murray Darling Basin, including the catchments of the Maranoa and Balonne-Culgoa river systems.

Major Primary Industries

The region has rich cropping and grazing lands and significant reserves of coal seam gas, conventional gas and petroleum. The agricultural economy includes irrigated horticulture and cotton, dryland cropping, intensive livestock (dairy), forestry, and grazing. The gross value production (GVP) in 2014-15 of agricultural commodities in the Border Rivers and Maranoa Balonne regions were \$1.5 B or 13% of the state total GVP for agricultural commodities (\$11.9 B, ABS 2016).



Sunset on the Balonne River, St George, Queensland

Courtesy of Tourism Queensland

Climate Trends and Projections

Historical changes in the key climate variables relevant to agricultural production including temperature, evaporation and rainfall are summarised in Table 1. Table 2 provides information on the historical means for the key variables and the projected changes for 2030.

Table 1: Historical Climate Trends (Interpreted and summarised from BoM 2016)

Variable	Trend Since (year)	Change per decade		
		Annual	Summer	Winter
Maximum Temperature (°C)	1950	+0.15 to +0.40	+0.05 to +0.20	+0.15 to +0.40
Minimum Temperature (°C)	1950	+0.10 to +0.40	+0.05 (east) to +0.30	+0.10 to +0.30
Mean Temperature (°C)	1950	+0.15 to +0.30	+0.15 to +0.30	+0.20 to +0.30
Pan Evaporation (mm)	1970	-10.0 (west) to +2.5	-5.0 to +2.5	-2.5 to NSC
Rainfall (mm)	1950	-30 (east) to 0 (west)	-10.0 to +5.0	-10 to NSC (south)

NSC - No significant change | Unknown of Hot Days, Warm Spell Duration or Growing Season Length | Pan Evaporation = the amount of water evaporated from an open pan per day

Additional climate projections for Queensland

- Global atmospheric **carbon dioxide concentration** (CO₂) is rapidly increasing. In March 2015, the monthly global average carbon dioxide concentration exceeded 400 ppm, well above the natural historical range from the last 800,000 years of 172 ppm to 300 ppm (CSIRO and BOM 2012a). Global CO₂ levels are projected to reach 540 ppm by 2050 and 936 ppm by 2100 (RCP8.5 high emissions) (IPCC 2013).
- Queensland can expect **longer dry periods** interrupted by **more intense rainfall** events. The frequency of both extreme El Niño and extreme La Niña events are likely to nearly double in response to greenhouse warming (Cai et al. 2014, 2015).
- The amount of time spent in **extreme drought** will increase in the highest emission scenarios (CSIRO and BOM 2015).

Table 2: Historical means for the period 1986-2005 and climate projections for 2030 (2020-2039) under the RCP8.5 emissions scenario relative to the model base period of 1986-2005

Variable		Annual	Summer	Autumn	Winter	Spring
Temperature (°C)	Historical mean	20.3	27.1	20.5	12.7	21.2
	Projections for 2030	+1 +0.2 to +1.7	+1 +0.3 to +2.0	+1 -0.1 to +1.7	+1 +0.4 to +1.7	+1 +0.2 to +1.9
Rainfall (mm)	Historical means	517	204	117	77	120
	Projections for 2030	-6% -18% to +6%	1% -19% to +23%	-6% -32% to +19%	-10% -45% to +18%	-7% -28% to +15%
Potential Evaporation (mm)	Historical mean	1598	<p style="text-align: center;">Historical means from 1986-2005</p> <p style="text-align: center;">Projections for 2030 (20-year period centred on 2030)</p> <p style="text-align: center;">Best Estimate</p> <p style="text-align: center;">Range of Change (5th - 95th)</p> <p style="text-align: center;"><i>For more information, including projections for 2050 and 2070, please refer to http://www.climatechangeinaustralia.gov.au/en/ or Ekström et al. 2015.</i></p>			
	Projections for 2030	+3% +1% to +7%				
Relative Humidity	Projections for 2030	-3% +5% to +6%				
Wind Speed	Projections for 2030	+1% -1% to +13%				



Grazing Cattle, Westmar, Queensland

Courtesy of Tourism Queensland

Impacts of a variable and changing climate in the Maranoa and Districts Region

Whilst a more variable and changing climate will impact the key primary industries in the region, the population and natural environment will also feel the effects.

Human Well-Being

The variable and changing climate of the region will have both direct and indirect impacts on health, location and living arrangements.

Likely Impacts	Potential Strategies for Adaptation
Extremes of weather and climate (drought, flood, cyclones, heatwaves etc.) on human well-being (TCI 2011, Hughes and McMichael 2011, NCCARF 2011a)	
<ul style="list-style-type: none"> • Direct effects of extremes of weather include injury and death during floods and cyclones, heat stress during heatwaves, and a reduction of cold-related deaths. • Indirect effects of extremes of weather could include an increase in the: <ul style="list-style-type: none"> ○ number of bushfires due to extreme heat and aridity; ○ risk of mosquito-borne, water-borne and food-borne diseases; ○ number of infectious and contagious diseases with an increase in the number of injuries; and ○ incidence of disease from microbial food poisoning with an increase in temperature. • Increases in extreme events can lead to increased pressure on health systems, including an increased demand for health professionals, ambulance and hospital workers. • Rural, regional and remote communities are particularly exposed in a changing climate compounding the chronic difficulties and inequities that already face many communities. Many parts of the country already find it hard to recruit dedicated health care and social service professionals. A changing climate will also increase the demand for social support and mental health services, and, at the same time, make it harder to recruit and retain staff in affected areas. • Severe weather events can destroy places and disrupt livelihoods and communities leading to long-term mental health effects. According to Bonanno et al. (2010), a significant part of the community, as many as one in five, will suffer the de-bilitating effects of extreme stress, emotional injury and de-spair. • The emotional and psychological toll of disasters can linger for months, even years, affecting whole families, the capacity for people to work and the wellbeing of the community. • Evidence is beginning to emerge that drought and heatwaves lead to higher rates (by about 8%) of self-harm and suicide (Doherty and Clayton 2011). • Those most vulnerable to extremes of weather and climate include children, the elderly, Indigenous communities and people with pre-existing diseases and disabilities. 	<ul style="list-style-type: none"> • Adapt existing buildings and plan any new infrastructure to take into account climate impacts and extreme events such as flooding. • Implement control measures to reduce the impact of bushfires, heatwaves, mosquitoes, water-borne and food-borne diseases, infectious and contagious diseases and injuries. • Continue to obtain information on the expected effects of a changing climate. • Develop agreements with your workers on how to manage extreme hot days, or identify periods of time where weather and climate affect working conditions. • Develop social support networks. • Contact your local council or relevant government department to find information on social and health support programs.

Biodiversity

The Brigalow Belt (BB) and Mulga Lands (ML) bioregions are present within the Maranoa and Districts region. The BB is the largest bioregion in Queensland and is very rich in species, including large numbers of plants and animals with small ranges. This bioregion has endemic and near-endemic eucalypt, wattle and invertebrate species. Dominant plants, including Mulga (*Acacia aneura*), on the ML bioregion are very resilient to a variable and changing climate, although deaths due to drought may still occur. The degree of ecological change caused by climate change is more likely to be greater in the plant biological group than that of mammals, amphibians or reptiles (Williams et al. 2014).

Likely Impacts	Potential Strategies for Adaptation
Extremes of weather and climate (drought, flood, cyclones, heatwaves etc.) on Biodiversity (Low 2011)	
<p>Impacts in the Brigalow Belt</p> <ul style="list-style-type: none"> Severe drought in the BB may result in deaths of many trees including Brigalow and Cypress pines. Buffel grass invasion is of particular concern within the BB. Invasion of this species may displace groundcover plants and significantly increase fire risk. <p>Impacts in the Mulga Lands</p> <ul style="list-style-type: none"> Within the ML, mass deaths of birds, mammals, reptiles and amphibians may occur if a heatwave of unprecedented severity struck during a severe drought. Koalas and yellow-footed rock wallabies seem most likely to decline because they are at the edges of their ranges within the ML. Impacts from reduced rainfall and increased temperature within the ML will worsen the impacts of invasive species, especially goats and buffel grass. 	<ul style="list-style-type: none"> Fire management. Manage weeds and invasive pasture grasses, such as buffel grass and guinea grass, to prevent spread into conservation areas and the habitats of rare species. Control pests and feral animals (goats, horses) to reduce losses and protect rare plants. Protect refugia habitats. Prevent vegetation thickening in conservation areas.

Grazing Industry

Cattle, sheep and wool are important primary industries in Queensland. In 2014-15 their combined GVP was \$5.2 B (44% of the total Queensland GVP of agricultural commodities, ABS 2016a) which is made up of the production and marketing of beef cattle (\$5.1 B), sheep and lambs (\$66.4 M) and wool (\$66.2 M).

Cattle numbers in Border Rivers and Maranoa Balonne (BRMB) regions were 1.2 M 2014-15 which was 10% of the total cattle numbers for Queensland (ABS 2014-15). In 2014-15 the GVP for cattle, sheep and wool for BRMB was \$570 M (ABS 2016a) or 5% of state and 37% of the value of BRMB agricultural commodities.

The majority of beef, sheep and wool production come from native pastures which cover about 85% of Queensland. The main pasture communities in MD are *Aristida-Bothriochloa* (56% of region), Brigalow (22%) and Mitchell grass (10%) (Tothill and Gillies 1992). The soil fertility is excellent (Brigalow) to poor (*Aristida-Bothriochloa*) and growth of pastures is usually limited by inadequate rainfall.

Case Study - Impacts in the Maranoa and Districts Region

The impacts of a changing climate are complex because of interacting and opposing forces operating within the biophysical system (McKeon et al. 2009). The process of assessing the impacts of a changing climate often involves deriving the 'best estimate' projections of future climate, simulating the grass growth and grazing strategies under changing climate conditions using well-calibrated grass/grazing system models, and combing the simulation output with successful producer and researcher experience in regional Queensland. A good example of a proven process of assessing the impacts, adaptive responses, risks and vulnerability associated with a changing climate is the 'risk matrix' approach (<http://www.longpaddock.qld.gov.au/products/matrix/index.html>, Cobon et al. 2009, 2016) which is customised for primary industries and is based on the Australian and New Zealand Risk Management Standards (Standards Australia 2004).

There are many gaps in knowledge, for example, the future climate projections are uncertain (particularly for rainfall) and in some cases the projected changes in rainfall and temperature appear smaller than to year-to-year variability. Nonetheless, a risk-averse approach to grazing management based on the 'best estimate' projections in combination with short-term management of climate variability is likely to take advantage of any opportunities and reduce the risk of adverse impacts. There are major known uncertainties in identifying the impacts of a changing climate in the grazing industry in relation to:

- 1) carbon dioxide and temperature effects on pasture growth, pasture quality, nutrient cycling and competition between grass, trees and scrubs;
- 2) the future role of woody plants including the effects of fire, climatic extremes and management of stored carbon (see

McKeon et al. 2009 for more detail); and

- 3) carbon dioxide effects on diet quality and liveweight gain of cattle (Stokes 2011).

Modelling analyses of native pasture grasses (C4 tropical and sub-tropical grasses) for the MD region were undertaken for the St George, Mitchell and Miles areas (Cobon et al. 2012 *unpublished data*, Table 3). The average impacts of future climate scenarios from the three locations were examined for pasture growth, pasture quality (% nitrogen of growth), liveweight gain of cattle (LWG kg/ha), frequency of burning and frequency of green pasture growing days (GPGD). The baseline climate period was 1960-1990 and carbon dioxide concentration was 350 ppm. Improvements in water and nitrogen use efficiency resulting from doubling of carbon dioxide levels were accounted for in the modelling as per Stokes 2011. The impacts were either positive or negative, and as a guide were also classified as being of either High (>20% change from baseline, H), Medium (5%-20%, M) or of little or no impact (5 to -5%, LC). The soils were of average fertility (20 kgN/ha) and the density of trees (7.11 m²/ha tree basal area) resembled that of open woodland.

Table 3: Matrix showing potential opportunities and risks associated with the average impacts of future climate scenarios from St George, Mitchell and Miles for modelled pasture growth (kg/ha), pasture quality (% nitrogen in growth), liveweight gain of cattle (LWG kg/ha), frequency of burning and green pasture growing days (GPGD) (Source: Cobon et al. 2012 *unpublished data*).

Future climate	Growth	Quality	LWG	Burning	GPGD
+3°C	LC	LC	+M	-M	+M
2xCO ₂	+H	-M	+H	LC	LC
+3°C, 2xCO ₂	+H	-M	+H	+H	+M
+3°C, 2xCO ₂ , +10% rainfall	+H	-H	+H	+H	+M
+3°C, 2xCO ₂ , -10% rainfall	+M	-M	+M	LC	+M

H= high, M= medium, LC = little change
 Shading indicates positive and negative impacts
 Positive impacts showing either High or Medium opportunities
 Negative impacts showing either High or Medium risks

This study found that there are likely to be:

- the future climate studied here is likely to have a positive impact on pasture growth, liveweight gain, frequency of burning and green pasture growing days, and a negative impact on pasture quality; and
- a 3°C rise in temperature is likely to reduce the frequency of burning, however, this may be outweighed by the positive impacts of a combined 3°C rise in temperature and doubled carbon dioxide (providing more opportunity for prescribed burning to control weeds, regrowth and dry vegetation).

Opportunities for the Grazing Industry

- Increased production of biomass from rising carbon dioxide levels as plants use water, nutrients and light resources more efficiently (Nowak et al. 2004).
- Improved plant water use efficiency will allow pastures to produce more biomass using the same amount of water (Stokes et al. 2011).
- Elevated carbon dioxide will increase the efficiency of water and nitrogen use by the pastures (Stokes et al. 2008), but this increase in growth of pastures is likely to be offset by a reduction in overall pasture quality (lower protein and lower digestibility) (Stokes et al. 2011).

Case Study – Impacts on pasture communities in the Maranoa

Modelling studies in the Maranoa on the combined effects of elevated carbon dioxide, a 2.3°C rise in temperature and 7% lower rainfall on three land types with different fertility showed the moderate fertility land types (e.g. Poplar box flats) to be more adversely affected compared to the more (e.g. Poplar box with buffel) or less fertile (e.g. Narrow leafed iron bark) land types. For example, Poplar box flats showed a 22% decline in forage production and a 55% reduction in carrying capacity (Phelps 2011).

Case Study - Using past records to help understand future impacts

Projected changes in rainfall of the order of ±10% appear low compared to year-to-year variability, or even in the difference between the average of El Niño and La Niña years (-20% and 20% rainfall respectively in eastern Australia) (McKeon et al. 2004). However, when the historical range of variation is analysed for a 25-year (climate change time-scale) moving average then a change in rainfall of ±10% is relatively high. For example, the 25-year moving average of rainfall at St George has fluctuated between -18 and +15% compared with the long-term average since 1890 (Figure 1). The extended periods of lower rainfall (1920s to 1950s, 1990s to 2000s) have been associated with extensive droughts, degradation events, reduced profits and greater debt and human hardship. It is likely that under drier climatic conditions these circumstances will become more familiar with shorter and less frequent recovery periods.




Figure 1: 25-year moving average rainfall (12 months, April in year 1 to March in year 2) at St George, Qld (Source: Clewett et al. 2003).

Likely Impacts	Potential Strategies for Adaptation
<p>Changed rainfall patterns</p> <ul style="list-style-type: none"> • Longer and more frequent droughts associated with more extremes of climate, fewer recovery events, changes in decadal rainfall variability and ENSO will decrease forage production, surface cover, livestock carrying capacity, animal production and cause major changes in plant and animal species composition (Cobon et al. 2009, McKeon et al. 2009). • Erosion risks are likely to increase due to greater year-to-year variability in rainfall. • Rising tree densities and declining pasture condition raise the sensitivity of pastures to climate induced water stress. 	<ul style="list-style-type: none"> • Manage perennial grass cover using ‘best management practice’ for the pasture community. For example, set the annual stocking rate at the end of each growing season to utilise a safe proportion (10-20%) of available pasture and make adjustments accordingly for beneficial or spoiling rainfall in winter or spring, early breaks to the dry season, locust plagues and forecasts of rainfall for the coming summer. • Monitor trends in rainfall. • Use climate indicators to make early adjustments in animal numbers. • Manage non-domestic grazing pressure. • Use wet season spelling of pastures. • Manage invasive plant species. • Maintain refugia especially around wetlands (Cobon et al. 2009). • Manage climate variability and change by using forecasts of rainfall (and temperature) in decision making. • Manage intra-seasonal (MJO, 30-60 day cycle), inter-annual (ENSO, 2-7 year cycle) and decadal rainfall variability (PDO/IPO, 20-30 year cycle) using indicators of MJO, ENSO (SOI, SST) and PDO, and climate analysis tools to adjust animal numbers commensurate with past and projected climate trends, such as: <ul style="list-style-type: none"> ○ LongPaddock (http://www.longpaddock.qld.gov.au); ○ AussieGRASS (http://www.longpaddock.qld.gov.au/about/researchprojects/aussiegrass/index.html); ○ ClimateArm http://www.armonline.com.au/ClimateArm ○ Bureau of Meteorology Website http://www.bom.gov.au, http://reg.bom.gov.au/climate/mjo; • Use supplementary feeding, early weaning and culling animals at risk to reduce mortalities in dry conditions (Fordyce et al. 1990). • Increase or maintain <i>Bos indicus</i> content in herd to increase cattle tick and buffalo fly resistance/resilience. • Monitor spread of pests, weeds and disease. • Introduce more species of dung fauna (control of buffalo fly larvae). • Promote greater use of traps and baits (buffalo and sheep blowflies) and vaccines (cattle ticks and worms). • Use fire to control woody thickening.

Likely Impacts	Potential Strategies for Adaptation
<p>Increased temperatures</p> <ul style="list-style-type: none"> Warming will be greatest toward the interior of the continent away from the moderating influence of the ocean. Each 1°C increase in temperature will cause a warming that would be roughly equivalent to moving about 145 km (or about 2° in latitude) closer to the equator (Stokes et al. 2011). For example, Roma under warming of 3°C is likely to receive temperatures currently experienced at Pentland (Figure 2). Livestock will be exposed to a greater risk of heat stress particularly in open grasslands. They are unlikely to travel as far to water which concentrates grazing pressure and increases the risk of adverse pasture composition changes and soil degradation (Howden et al. 2008). Increased day time temperatures increases water turn-over and evaporative heat loss resulting in reduced rate of passage and forage intake in livestock (Daly 1984). Increased night time temperatures can reduce recovery time of livestock and increase the effects of heat stress during the day. Increased heat stress reduces fertility, conception, peri-partum survival and follicle development in sheep. Warmer conditions favour vectors and the spread of animal disease (White et al. 2003). Pastures could cure earlier under warmer climates shifting the timing of fires to earlier in the season. Warmer drier conditions with higher frequency of storms could increase the risk of wildfires. 	<ul style="list-style-type: none"> Arrange water points to reduce distance to water and even out grazing pressure. Select the time of mating to optimise nutritional requirements and reduce the risk of mortality in new-borns. Select cattle lines with effective thermoregulatory controls, efficient feed conversion and lighter coat colour (Finch et al. 1984, King 1983). Proactively control disease by targeting known sources of disease and vectors (Sutherst 1990). Maintain high standards of animal welfare to build domestic and export meat and fibre markets (Mott and Edwards 1992). Incorporate greater use of prescribed burning to reduce the risk of wildfires and control woody thickening. Rotate paddocks of heavier grazing for use as fire breaks. Maintain or improve quarantine capabilities, monitoring programs and commitment to identification and management of pests, disease and weed threats. Develop species resistant to pests and disease, and use area-wide improved management practices.




Figure 2: Annual average temperature in Australia (Source: Bureau of Meteorology). One degree of warming is roughly equivalent to moving 145 km toward the equator.

Likely Impacts	Potential Strategies for Adaptation
Increased temperature, higher carbon dioxide concentration and changed rainfall	
<ul style="list-style-type: none"> • Pastures growing under a climate characterised by consistent water stress appear to benefit most from increased plant water use efficiency under elevated carbon dioxide. • The fertilisation effects of doubled carbon dioxide (700 ppm) were found to offset declines in forage production under 2°C warming and a 7% decline in rainfall (Webb et al. 2011). • The combined effects of elevated carbon dioxide (650 ppm), higher temperature (3°C) and lower rainfall (10%) resulted in 10-20% lower forage production (McKeon et al. 2009). In this study increased temperature and declining rainfall outweigh the conservatively represented benefits of increasing carbon dioxide. • Rising carbon dioxide will result in a reduction in overall pasture quality (lower protein and lower digestibility) (Stokes et al 2011). 	<ul style="list-style-type: none"> • Maintain land in good condition to reduce potential declines in forage production under a warmer drier climate. • To compensate for declining forage quality, increase the use of supplements (N, P and energy) and rumen modifiers. • Destock earlier in the season to make greater use of feedlots to finish livestock. • Explore alternative land use in marginal areas. • Apply safe carrying capacity of ~10-15% utilisation of average long-term annual pasture growth. • Undertake risk assessments to evaluate needs and opportunities for changing species, management of land and land use. • Support assessments of the benefits and costs of diversifying property enterprises. • Introduce pasture legumes to improve nitrogen status.
More intense storms	
<ul style="list-style-type: none"> • Rainfall intensity is expected to increase as temperature and moisture content of the atmosphere increase. • A 1°C increase in temperature may result in an increase in rainfall intensity of 3-10% (SAG 2010). • More intense storms are likely to increase runoff, reduce infiltration, reduce soil moisture levels and pasture growth, and increase the risk of soil erosion. 	<ul style="list-style-type: none"> • Maintain pasture cover for optimal infiltration of rainfall. • Adjust livestock numbers to maintain good coverage of perennial pastures during the storm season.
Higher temperature humidity index (combination of maximum temperature and dewpoint temperature)	
<ul style="list-style-type: none"> • Temperature humidity index (THI) is an indicator of heat stress. Heat stress in beef cattle is significant at a THI of over 80. Frequency of days per year above this level is shown in Figure 3 for historical and projected climate. Rising temperature by 2.7°C increases the occurrence of heat stress by about 30% points (Howden et al. 1999). • Heat stress reduces liveweight gain and reproductive performance in beef cattle, and increases mortality rates (see Howden et al. 1999). • Heat stress reduces the development of secondary wool follicles in sheep, reducing lifetime wool production in sheep (Hopkins et al. 1978). 	<ul style="list-style-type: none"> • Select cattle lines with effective thermoregulatory controls, efficient feed conversion and lighter coat colour (Finch et al. 1984, King 1983).




Figure 3: Frequency of days per year that the THI>80 for a) 1957-97 and b) a future climate scenario of +2.7°C. Thermal stress is significant in beef cattle when the THI exceeds 80 (Source: Howden et al. 1999).

Dairy Industry

In 2014-15 the Queensland dairy industry had a herd of about 168,000 dairy cattle of which 91,000 are cows in milk (ABS 2016b). The Queensland dairy industry produced 411 million litres of milk from 448 farms, which was 4.2% of Australia's milk production (Dairy Australia 2015). In 2014-15 Border Rivers and Maranoa Balonne region produced 2% of the value of Queensland's whole milk (\$236 M, ABS 2016a).

Much of the information below on the impacts of a changing climate for the dairy industry is drawn from Dairy Australia (2011).

Opportunities for the Dairy Industry

- Increased plant photosynthesis and associated increased production with increases in carbon dioxide.
- Increased pasture growth during cooler months due to increased minimum temperatures and less frosts.
- Lower water availability will favour short rotation pasture systems.

Case Study - The effects of increased temperature on dairy cows.

Cows have the ability to off-load heat; however prolonged periods of heat, particularly above 25°C, may lead to heat stress. Heat stress reduces the cows' ability to produce milk and get in calf. There may also be health and welfare problems.

Management and adaptation tools to minimise the risk of heat stress include increased provisions of shade, active cooling sprays and breed selection.

Likely Impacts	Potential Strategies for Adaptation
Increased temperatures	
<ul style="list-style-type: none"> • Rising temperatures may cause an increase in the incidence of heat stress to dairy cows. • Higher temperatures may make C4 pasture species more competitive at the expense of nutritious C3 species, however higher carbon dioxide is expected to favour C3 species more than C4. • Water and irrigation requirements may be increased with higher temperatures. 	<ul style="list-style-type: none"> • Provide more cooling mechanisms for dairy cows e.g. shade and active cooling sprays. • Selectively breed stock, pasture and feedstock for their ability to withstand higher temperatures. • Switch to pasture species that will adapt to changing conditions. • Sow pastures earlier to match warmer conditions. • Use nitrogen fertiliser during winter months. • Use short rotation pasture systems and winter fodder crops.
Decreased rainfall	
<ul style="list-style-type: none"> • There may be associated lower runoff and reduced soil moisture. • Less water will be available causing more competition for water. • Lower growth of rain-fed pastures and crops. 	<ul style="list-style-type: none"> • Decrease evaporation rates in water storage and in the soil. • Install more efficient irrigation systems and improve water use efficiency. • Change feed system. • Apply more emphasis to crops. • Switch to pasture species that will adapt to changing conditions.
More intense and frequent storms with increased seasonal variability	
<ul style="list-style-type: none"> • Livestock could be injured by more intense storms and hail, particularly in intensive production systems where animals are concentrated. • Extreme wet seasons can negatively impact milk production, herd health and property infrastructure. 	<ul style="list-style-type: none"> • Use summer housing for dairy cows. • Develop and implement a risk management plan when long range weather forecasts indicate a higher than average probability for either a wet or dry season ahead.

Cropping Industry

Broadacre cropping in Queensland produces a range of cereal, oilseed and legume crops, including wheat, maize, barley, sorghum, chickpea, mungbean, soybean, sunflowers and peanuts (QFF 2012). In Queensland the most commonly grown winter crop is wheat (1 M tonnes in 2014-15, ABS 2016b) and summer crop is sorghum (1.6 M tonnes in 2009-10, ABS 2016b). In 2014-15 the value of broadacre crops, excluding crops harvested for hay, cotton and sugar was \$1.1 B (ABS 2016a) in Queensland and \$365 M in Border Rivers and Maranoa Balonne (BRMB) (ABS 2016a). In 2014-15 the value of cotton was \$175 M (46% of state GDP) and pasture and cereal crops cut for hay was \$13 M in BRMB (ABS 2016a).

Broadacre cropping across the region will be affected by climate change at both the enterprise scale and regionally. The information below details the impacts of a changing climate on the cropping industry and adaptation responses that may result in a more resilient system (Cobon et al. 2016, Stokes and Howden 2010 and references therein).

Opportunities for the Cropping Industry

- Increased carbon dioxide may result in higher crop yields and biomass due to increased carbon dioxide fertilisation and photosynthesis.
- C3 plants (cereal grain crops like wheat) respond better to increased carbon dioxide than C4 plants (tropical-origin crops such as sugar cane and maize).
- The effect of increased temperature may, however, have the opposite effect due to increased water stress. Therefore the net results remain uncertain (NCCARF 2011b).
- In cooler months, increased temperatures may reduce frost risk.

Case Study – Effects of higher temperatures and increased carbon dioxide on the growth of cotton

Simulation modelling of cotton growth indicates that an increase in temperature and a 20% decrease in rainfall would cause a decrease in cotton yield of 20%. However, an increase of atmospheric CO₂ (to 555 ppm) will moderate such effects to only a 4% decrease in yield (Williams et al. 2015).

Likely Impacts	Potential Strategies for Adaptation
Increased temperatures and carbon dioxide concentration	
<ul style="list-style-type: none"> • Rising carbon dioxide may increase biomass production and grain yields which will in turn reduce both the average nitrogen level of grain and the frequency of achieving key nitrogen thresholds. • Warmer temperatures and increased rainfall are likely to favour the slower-maturing cultivars (greater thermal time requirements) that could benefit from an earlier date of flowering and a longer period of photosynthesis (with adequate moisture). • Heat stress during the summer months is likely to cause poor seed set in summer grain crops, such as mung bean, sunflower and maize because higher temperatures lead to earlier flowering crops and poor pollination. • Heat stress during spring may decrease yield of winter crops (e.g. wheat). • Warmer temperatures in spring may allow earlier planting of summer crops with lower frost risk. • Decreased frost incidence may benefit winter crops because of less chance of frost at flowering, however this will be complicated by the fact that they will flower earlier. 	<ul style="list-style-type: none"> • Adjust planting times of summer crops (e.g. mung beans, sunflower and maize) so that they are not flowering during the hottest months. • To maintain grain nitrogen content at historical levels, there will be a need to increase fertiliser application rates by up to 50% depending on the yield expectations. Therefore, increase nitrogenous fertiliser application or increase use of pasture legume rotations may be needed to maintain grain yields and protein content. • Increase application rates of other crop nutrients (e.g. P, K).
Changed rainfall patterns and increased storm frequency	
<ul style="list-style-type: none"> • Increased risk of storm damage and erosion. • Increased occurrence of some pests and diseases. • Heavy rainfall can increase leaching of nutrients and movement of salts, although total rainfall is likely to decline. • Decreased yields as a result of increased crop water stress. 	<ul style="list-style-type: none"> • Optimise availability of all resources (e.g. through precision agriculture). • Adopt efficient irrigation technology to control water table, monitor water table position and improve catchment vegetation distribution and ground cover to increase infiltration rate. • Apply fungicides to wheat crops to decrease leaf disease (Meinke and Hochman 2000). • Reduce soil moisture loss by: <ul style="list-style-type: none"> ◦ increasing residue cover by minimal or no-tillage; ◦ establishing crop cover in high loss periods; ◦ weed control; and ◦ maximising capture and storage of excess rainfall on-farm. • Use cover crops to increase groundcover during long fallows or after crops that don't produce much cover. • Establish a higher percentage of summer crops relative to winter crops as rainfall changes point towards the largest decreases being in winter and spring. • In mixed farming systems, where cropping is marginal and may become more so, consider incorporating a greater proportion of livestock into the farm business for profitability.

Likely Impacts	Potential Strategies for Adaptation
Increased temperatures and decreased rainfall	
<ul style="list-style-type: none"> • Warmer temperatures and a significant decrease in rainfall are likely to favour winter crop varieties (e.g. wheat and barley) with earlier-flowering characteristics which allow grain-fill to occur in the cooler, wetter parts of the year in dry areas. Varieties with characteristics such as higher response to elevated carbon dioxide conditions, rapid germination, early vigour and increased grain set in hot/windy conditions may also be favoured. • Increased temperatures and evaporation may reduce the yield of dryland crops like wheat and sorghum (Potgieter et al. 2004); however, this may be offset by increased carbon dioxide. • Irrigated crops may be adversely affected due to a reduction in supply of irrigation water. • There will be more pressure and challenges for managing groundcover, crop choice (winter or summer), soil nutrient requirements, pest and weed control, soil carbon etc., especially from higher temperature, increased soil moisture stress and higher rainfall variability. • Lower rainfall may reduce deep drainage in dryland cropping systems. 	<ul style="list-style-type: none"> • Incorporate 'best practice' farm management by constantly varying crops and inputs based on the availability of limited and variable resources and signals from the operating environment (Rodriguez et al. 2011a, Rodriguez et al. 2011b). • Use varieties that incorporate the traits of appropriate thermal time (degree days) and vernalisation (exposure to cold temperatures required for flowering) requirements and with increased resistance to heat shock and drought. • Diversify the farm enterprise (e.g. using opportunistic planting). • Increase the use of legume-based pastures and leguminous crops or further increase nitrogen fertiliser application to maintain grain quality, especially protein content. • Adjust planting times to cater for changes in crop maturity and the duration and timing of heatwaves. • Adopt efficient irrigation technology. • Increase use of supplementary water. • Optimise irrigation scheduling. • Use more effective irrigation water delivery technologies (i.e. trickle tape). • Construct on-farm water storage facilities. • Use drought-tolerant or more water efficient varieties. • Modify row spacing. • Minimise tillage. • Use cover crops. • Manage water resources and improve efficiency of irrigation systems. • Integrate cropping into regions of higher rainfall. • Make crop planting decisions based on seasonal climate forecasting, soil tests and other climate related information obtained from tools such as Rainman, Whopper Cropper and APSIM. • Use adaptive crop management techniques such as: <ul style="list-style-type: none"> ○ zero-tillage practices, minimum disturbance planting techniques (e.g. seed pushing); ○ controlled traffic; ○ responding to planting opportunities when they occur; ○ widening row spacing or skip-row planting; ○ lowering plant populations; ○ using efficient on-farm irrigation management with effective scheduling, application and transfer systems; and ○ assessing fertiliser inputs. • Reduce surface soil erosion by: <ul style="list-style-type: none"> ○ increasing residue retention; ○ maintaining erosion control infrastructure (e.g. contour banking); and ○ using controlled traffic systems. • Control pests and diseases.

Horticulture Industry

Horticulture is Queensland's second largest primary industry (QFF 2012). Queensland grows approximately one third of Australia's horticulture produce, with more than 120 different types of fruit and vegetables being grown in 16 defined regions covering a total area of 100,000 hectares and 2800 farms (QFF 2012, HAL 2012). In 2014-15 the value of production for Queensland was about \$2.5 B which was made up of \$1 B for vegetables, \$1.2 B for fruit and nuts and \$290 M for nurseries, cut flowers and turf (ABS 2016a).

In 2014-15 the Border Rivers and Maranoa Balonne (BRMB) produced about 9% of the total value of the state's horticulture, including 7% of the value of vegetables, 7% of the value of fruit and nuts, and 9% of the value of nurseries, cut flowers and turf (ABS 2016a). The BRMB is a major producer of Queensland's pome fruit, stone fruit and grapes.

Much of the information below on the impacts of a changing climate for the horticulture industry is drawn from reports commissioned for the Garnaut Review (Deuter 2008).

Opportunities for the Horticulture Industry

- Increased minimum temperature, reduced frost frequency and shortened frost period during the growing season may alter production windows.
- Vegetable growers will have the additional option of planting later, therefore extending the production season.

Case Study – The effects of increased temperatures on Pumpkin crops

Pumpkins are known to be a warm season crop; however, prolonged hot weather causes poor fruit set. They are very sensitive to frost with all stages being affected by cool temperatures. Therefore an increase in minimum temperatures may be beneficial to the pumpkin growing industry (Lovatt 2010).



Likely Impacts	Potential Strategies for Adaptation
Increased temperatures	
<ul style="list-style-type: none"> • Changes to the suitability and adaptability of some crops. • Potential shift in the optimum growing regions from the current hotter producing areas towards areas currently regarded as too cool. • Change the timing and reliability of plant growth, flowering, fruit growth, fruit setting, ripening and product quality; fruit size, quality and pollination. • Change harvesting times for different areas. • Reduce the time to reach maturity (earlier in the season). • Change the occurrence and distribution patterns of fruit fly and heliothis • Potentially downgrading product quality. • Result in pollination failures. • Increase active soil-borne diseases and insect infestation for longer periods during the year. • May cause fruit quality issues to be more common early and late in the season and may influence fruit quality and pollination. • Reduced diurnal temperature range will potentially reduce the overlap between open stages of male and female flower parts thus decreasing the chances for pollination and resulting in more pollination failures, fruit drop and sunburn to fruit. • Increased minimum temperatures and reduced occurrence of frost may benefit some production systems if managed correctly. The winter production season will be shortened. • Changes in disease and pest distribution ranges. 	<ul style="list-style-type: none"> • Select for, or change to, cultivars which are more adaptable to a changing and variable climate. • Select and review growing site/location to avoid unsuitable climate factors through identifying threshold temperatures or other climate conditions for crops. • Choose optimal timing of planting. • Start breeding programs for heat tolerant, low chill, and more adaptable varieties of various horticultural crops. Varieties with higher quality under enhanced carbon dioxide and elevated temperatures will need to be evaluated then considered in breeding programs. • Apply the latest research results and best management techniques to maintain product quality. • Use crop protection treatments including solar radiation shading and evaporative cooling through overhead irrigation to maintain fruit quality. • Use tools/models associated with managing climate variability to improve both quality and quantity of horticulture products. • Plant varieties with chilling requirements below 1000 hours.
Changed rainfall patterns	
<ul style="list-style-type: none"> • Increased risk to crops reliant on irrigation where irrigation water availability is reduced especially during dry periods. • Changes to the reliability of irrigation supplies, through impacts on recharge to surface and groundwater storages. 	<ul style="list-style-type: none"> • Adopt more efficient irrigation monitoring and scheduling technologies which provide further water-use efficiencies. • Apply the latest research results and best management techniques to maintain product quality, including fertiliser timing and amounts according to crop requirements. • Use tools/models associated with managing climate variability to improve both quality and quantity of horticulture products.
More intense storms	
<ul style="list-style-type: none"> • Increased runoff may providing opportunities for growers to capture more water for irrigation. • Lead to conditions favouring foliar diseases and some root invading fungi, for example, the fungus <i>Phytophthora cinnamomi</i>. • Increase the likelihood of crop damage, decreasing quality and production. • Affect the timing of cultural practices and ability to harvest, as well as negative effects on yield and product quality. 	<ul style="list-style-type: none"> • Improve Integrated Pest and Disease Management (IPDM) practices to adapt to a changing climate and encourage disease suppressive soil techniques. • Improve on-farm water storage linked to drainage and water harvesting systems. • Improve sediment runoff protection via grassed waterways and erosion control structures. • Improve plant nutrition management. • Improve all-weather access to cropping areas.



Dry Paddock, Roma, Queensland

Courtesy of Tourism Queensland

More Information

For more information, including projections for 2050 and 2070, please refer to <http://www.climatechangeinaustralia.gov.au/en/> or Ekström et al. 2015.

For more information on the varying and changing climate please see the Queensland Government and The Long Paddock websites at <http://www.qld.gov.au/environment/climate/climate-change/> and <http://www.longpaddock.qld.gov.au>, in particular:

- The Climate Change Risk Management Matrix - <http://www.longpaddock.qld.gov.au/products/matrix/index.html>
- Queensland Coastal Hazard Area Maps - http://ehp.qld.gov.au/coastal/management/coastal_plan_maps.php#map_layers

Citation

Cobon DH, Terwijn MJ, and Williams AAJ (2017). *Impacts and adaptation strategies for a variable and changing climate in the MARANOVA AND DISTRICTS REGION*. International Centre for Applied Climate Sciences, University of Southern Queensland, Toowoomba, Queensland, Australia.

Acronyms

APSIM, Agriculture Production Simulation Model
 ENSO, El Niño Southern Oscillation
 IPO, Interdecadal Pacific Oscillation
 GVP, Gross Value of Production
 MJO, Madden Julian Oscillation or 40 day wave
 PDO, Pacific Decadal Oscillation
 SOI, Southern Oscillation Index
 SST, Sea Surface Temperature

The authors would like to acknowledge the expertise in reviewing this document by staff in the Queensland Department of Science, Information Technology and Innovation, the Queensland Department of Agriculture and Fisheries, CSIRO and Dairying for Tomorrow.

References

- Australian Bureau of Statistics (ABS) 2016a, Value of agricultural commodities produced, Australia, 2014-15, Category Number: 7503.0, Australian Bureau of Statistics © Commonwealth of Australia, <http://www.abs.gov.au>
- Australian Bureau of Statistics (ABS) 2016b, Agricultural Commodities, Australia, 2014-15, Category Number: 71210.0, Australian Bureau of Statistics © Commonwealth of Australia, <http://www.abs.gov.au>
- Bonanno, GA, Brewin, CR, Kaniasty, K and La Greca, AM 2010, Weighing the Costs of Disaster: Consequences, Risk, and Resilience in Individuals, Families, and Communities, *Psychological Science in the Public Interest*, vol. 11, pp. 1-49.
- Booth, D, Edgar, G, Figueria, W, Jenkins, G, Kingsford, M, Lenanton, R and Thresher, R 2009, Temperature Coastal and Demersal Fish and Climate Change, A Marine Climate Change Impacts and Adaptation Report Card for Australia 2009.
- Bureau of Meteorology (BoM) 2016, Australian climate variability and change - Trend Maps, Bureau of Meteorology, <http://reg.bom.gov.au/cgi-bin/climate/change/trendmaps.cgi>
- Cai, W., Borlace, S., Lengaigne, M., Van Rensch, P., Collins, M., Vecchi, G., Timmermann, A., Santoso, A., McPhaden, M.J., Wu, L. and England, M.H. 2014, Increasing frequency of extreme El Niño events due to greenhouse warming. *Nature climate change*, 4(2), pp.111-116.
- Cai, W., Wang, G., Santoso, A., McPhaden, M.J., Wu, L., Jin, F.F., Timmermann, A., Collins, M., Vecchi, G., Lengaigne, M. and England, M.H. 2015. Increased frequency of extreme La Niña events under greenhouse warming. *Nature Climate Change*, 5(2), pp.132-137.
- Caldeira, K. and M. E. Wickett. 2005, "Ocean model predictions of chemistry changes from carbon dioxide emissions to the atmosphere and ocean." *Journal of Geophysical Research: Oceans* 110.C9.
- Canegrowers 2011, Annual Report 2010/11, Queensland Cane Growers Organisation Ltd, Brisbane, Queensland <http://www.canegrowers.com.au>
- Clewett, JF, Clarkson, NM, George, DA, Ooi, SH, Owens, DT, Partridge, IJ and Simpson GB 2003, Rainman StreamFlow version 4.3: a comprehensive climate and streamflow analysis package on CD to assess seasonal forecasts and mangement climate risk, Q103040, Department of Primary Industries, Queensland.
- Cobon, D, Stone, G, Carter, J, Scanlan, J, Toombs, N, Zhang, X, Willcocks, J and McKeon, G 2009, The Climate Change Risk Management Matrix for the grazing industry of northern Australia, *The Rangeland Journal*, vol. 31, pp.31-49 <http://www.longpaddock.qld.gov.au/about/publications/index.html>
- Cobon, DH, Williams, AA, Power, B, McRae, D and Davis, P, 2016. Risk matrix approach useful in adapting agriculture to climate change. *Climatic Change*, pp.1-17.
- Commonwealth Scientific and Industrial Research Organisation and Bureau of Meteorology (CSIRO and BOM) 2012a, State of the Climate 2012, Commonwealth Scientific and Industrial Research Organisation and Bureau of Meteorology <http://www.csiro.au/Outcomes/Climate/Understanding/State-of-the-Climite-2012.aspx>
- CSIRO and Bureau of Meteorology 2015, Climate Change in Australia Information for Australia's Natural Resource Management Regions: Technical Report, CSIRO and Bureau of Meteorology, Australia.
- Courtney, AJ, Kienzle, M, Pascoe, S, O'Neill, MF, Leigh, GM, Wang, Y-G, Innes, J, Landers, M, Braccini, M, Prosser, AJ, Baxter, P, Sterling, D & Larkin, J 2012, *Harvest strategy evaluations and co-management for the Moreton Bay Trawl Fishery*, Australian Seafood CRC final report, project 2009/774, Queensland Department of Agriculture, Fisheries and Forestry, Brisbane.
- Dairy Australia 2011, Climate Toolkit, Dairy Australia <http://www.dairyaustralia.com.au>.
- Dairy Australia 2015. Australian Dairy Industry in Focus, Dairy Australia Limited 2015. 2015ISSN 2202-7467 (Online); ISSN 1448-9392 (Print).
- Daly, JJ 1984, Cattle need shade trees, *Queensland Agriculture Journal*, vol. 110, pp.21-24.
- Deuter, P 2008, Defining the Impacts of Climate Change on Horticulture in Australia, Reports Commissioned by the Garnaut Climate Change Review, <http://www.garnautreview.org.au/CA25734E0016A131/pages/all-reports--resources-commissioned-reports.html>
- Doherty, TJ and Clayton, S 2011, The Psychological Impacts of Global Climate Change, *American Psychologist*, vol. 66, pp. 265-76
- Ekström, M, 2015. Central Slopes Cluster Report, Climate Change in Australia Projections for Australia's Natural Resource Management Regions: Cluster Reports, eds. Ekström, M. et al.
- Finch, VA, Bennett, IL and Holmes, CR 1984, Coat colour in cattle - effect on thermal balance, behaviour and growth, and relationship with coat type, *Journal of Agriculture Science*, vol. 102, pp. 141-47.
- Fordyce, G, Tyler, R and Anderson, VJ 1990, Effect of reproductive status, body condition and age of *Bos indicus* cross cows early in a drought on survival and subsequent reproductive performance, *Australian Journal of Experimental Agriculture*, vol. 30, pp. 315-22.
- Hopkins, PS, Knight, GI and Le Feuvre, AS 1978, Studies of the environment physiology of tropical Merinos, *Australian Journal of Agricultural Research*, vol. 29, pp. 161-71
- Horticulture for Tomorrow (HAL) 2012, Horticulture for Tomorrow: For Growers, Your Catchment, Queensland, http://www.horticulturefortomorrow.com.au/for_growers/your_catchments/qlld.asp.
- Hosking, C., Mills, M. and Lovelock, C. 2014, *Climate Change and Agriculture: a study for the SEQ Catchments NRM*, Global Change Institute, University of Queensland.
- Howden, SM, Crimp, SJ and Stokes, CJ 2008, Climate change and Australian livestock systems: impacts research and policy issues, *Australian Journal of Experimental Agriculture*, vol. 48, pp.780-88.
- Howden, SM, Hall, WB and Bruget, D 1999, Heat stress and beef cattle in Australian rangelands: recent trends and climate change, People and Rangelands, Building the Future, Proceedings of the 6th International Rangelands Conference, pp. 43-44.
- Hughes, L and McMichael, T 2011, The Critical Decade: Climate change and health, Climate Commission, Commonwealth of Australia (Department of Climate Change and Energy Efficiency), http://climatecommission.gov.au/wp-content/uploads/111129_FINAL-FOR-WEB.pdf
- Intergovernmental Panel on Climate Change (IPCC) 2013, *Scenario Process For Ar5*, http://sedac.ipcc-data.org/ddc/ar5_scenario_process/index.html
- King, JM 1983, Livestock water needs in pastoral Africa in relation to climate and forage, Addis Ababa, ILCA Research report, vol. 94.
- Lovatt, J 2010, Temperature requirements and limitations, Department of Primary Industries, Queensland Government, http://www.dpi.qld.gov.au/28_15331.htm
- Low, T 2011, Climate Change and Terrestrial Biodiversity in Queensland, Department of Environment and Resource Management, Queensland Government, Brisbane.
- McKeon, G, Hall, W, Henry, B, Stone, G and Watson, I 2004, Learning from history. Pasture degradation and recovery in Australia's rangelands, Queensland Department of Natural Resources, Mines and Energy, Brisbane.
- McKeon, GM, Stone, GS, Syktus, JJ, Carter, JO, Flood, NR, Ahrens, DG, Bruget, DN, Chilcott, CR, Cobon, DH, Cowley, RA, Crimp, SJ, Fraser, GW, Howden, SM, Johnston, PW, Ryan, JG, Stokes, CJ and Day, KA 2009, Climate change impacts on northern Australian rangeland livestock carrying capacity: a review of issues, *A Climate of Change in Australian Rangelands*, *The Rangeland Journal*, vol. 31, pp.1-30.
- Meat and Livestock Australia (MLA) 2010, Cattle numbers - as at June 2010: Natural Resource Management Region, Meat and Livestock Australia, <http://www.mla.com.au/About-the-red-meat-industry/Industry-overview/Cattle>.
- Meinke, H, and Hochman, Z 2000, Using seasonal climate forecasts to manage dryland crops in northern Australia—experiences from the 1997/98 seasons. In *Applications of Seasonal Climate Forecasting in Agricultural and Natural Ecosystems* (pp. 149-165). Springer Netherlands.
- Mott, JJ and Edwards, GP 1992, Beyond beef - a search for a level grazing paddock, *Search*, vol.7, pp. 223-5.
- Murphy, R and Simpson, G 2011, Balancing dairy production and profits in Northern Australia: QDAS Financial and production trends – 2011, Department of Employment, Economic Development and Innovation, Toowoomba, Queensland.
- National Climate Change Adaptation Research Facility (NCCARF) 2011a, Climate change impacts factsheet: 8. Human Health, <http://www.nccarf.edu.au/sites/default/files/8.Health-Impacts.pdf>
- National Climate Change Adaptation Research Facility (NCCARF) 2011b, Climate change impacts factsheet: 4. Primary Industries, <http://www.nccarf.edu.au/sites/default/files/4.Primary-Industries-Impacts.pdf>
- Nowak, RS, Ellsworth, DS and Smith, DS 2004, Functional responses of plants to elevated atmospheric CO₂ - do photosynthetic and productivity data from FACE experiments support early predictions? *New Phytologist*, vol. 162, pp. 253-80.
- Poloczanska, ES, Hobday, AJ and Richardson, AJ 2009, Report Card of Marine Climate Change for Australia. In: Poloczanska, ES, Hobday, AJ and Richardson, AJ (eds.), NCCARF Publication 05/09.
- Phelps, D 2011, Developing improved grazing and related practices to assist beef production enterprises across northern Australia adapt to a changing and more variable climate, B.NBP.0616 MLA Milestone Report 5, Meat and Livestock Australia.
- Potgieter, A, Meinke, H and Doherty, A 2004, Likely impact of climate on wheat and sorghum production in central Queensland, 4th International Crop Science Congress, Brisbane.
- Queensland Department of Agriculture, Fisheries and Forestry (QLDDAFF) 2010, Sugar, http://www.daff.qld.gov.au/26_6730.htm
- Queensland Farmer's Federation (QFF) 2012, Farming in Queensland, Queensland Farmer's Federation, <http://www.qff.org.au/farming-in-qlld/>

- Rodriguez, D, deVoil, P, Power, B and Cox, H** 2011a, Adapting to change: more realistic quantification of impacts and better informed adaptation alternatives, 5th World Congress of Conservation Agriculture incorporating 3rd Farming Systems Design Conference, September 2011, Brisbane.
- Rodriguez, D, deVoil, P, Power, B, Cox, H, Crimp, S and Meinke, H** 2011b, The intrinsic plasticity of farm businesses and their resilience to change. An Australian example, *Field Crops Research*, vol. 124, pp. 157-270.
- Scientific Advisory Group (**SAG**) 2010, Increasing Queensland's resilience to inland flooding in a changing climate, Final Scientific Advisory Group report, Department of Environment and Resource Management, Brisbane, <http://www.ehp.qld.gov.au/climatechange/pdf/inland-flood-study.pdf>
- Standards Australia** 2004, AS/NZS4360:2004 Risk Management.
- Stokes, C** 2011, Developing improved industry strategies and policies to assist beef enterprises across northern Australia adapt to a changing and more variable climate, MLA Milestone 3 report, Project No. B.NBP.0617, Meat and Livestock Australia.
- Stokes, C and Howden, M** 2010, Adapting agriculture to climate change: preparing Australian agriculture, forestry and fisheries for the future, CSIRO.
- Stokes, C, Ash, AJ, Scanlan, J and Webb, N** 2011, Strategies for adapting to climate change, Proceedings of the Northern Beef Research Conference, Darwin, pp. 81-86.
- Stokes, CJ, Ash, AJ and Holtum, JA** 2008, Savannas face the future: windows into a future CO₂ enriched world, A Climate of Change in the Rangelands, 15th Biennial Conference, Charters Towers, Queensland, Australian Rangeland Society.
- Sutherst, RW** 1990, Impact of climate change on pests and diseases in Australia, *Search*, vol. 21, pp. 232.
- The Climate Institute (**TCI**) 2011, A Climate of Suffering: the real cost of living with inaction on climate change (Melbourne and Sydney), The Climate Institute.
- Tothill, CJ and Gillies, C** 1992, The pasture lands of northern Australia - their condition, productivity and sustainability, Tropical Grassland Society of Australia, Occasional Publication No 5, Brisbane.
- Webb, NP, Stokes, CJ and Scanlan, JC** 2011, Interacting effects of vegetation, soils and management on the sensitivity of Australian savanna rangelands to climate change, *Climate Change*, DOI: 10.1007/s10584-011-0236-0.
- White, N, Sutherst, R, Hall, W and Whish-Wilson, P** 2003, The vulnerability of the Australian beef industry to the impacts of the cattle tick (*Boophilus microplus*) under climate change, *Climatic Change*, vol. 61, pp. 157-90.
- Williams, A, White, N, Mushtaq, S, Cockfield, G, Power, B and Kouadio, L**, 2015 Quantifying the response of cotton production in eastern Australia to climate change. *Climatic Change*, 129(1-2), pp.183-196.
- Williams KJ, Prober SM, Harwood TD, Doerr VAJ, Jeanneret T, Manion G, and Ferrier S** (2014) Implications of climate change for biodiversity: a community-level modelling approach, CSIRO Land and Water Flagship, Canberra. Available at: www.AdaptNRM.org ISBN 978-1-4863-0479-0.